





16700 Bayview Avenue, Unit #22, Newmarket, Ontario L3X 1W1
 Tel. 289-803-2959 • Fax. 1-855-870-1125
 1580-20th Street East, Unit 1, Owen Sound, Ontario N4K 5P5
 Tel. 519-372-9951 • Fax. 519-372-1384
 newmarketcardio@gmail.com
 191 Mcnaughton Rd., Suite 407, Maple, ON L6A 4E2
 Tel. 289-803-2959 • Fax. 1-855-870-1125
 www.advancedcardiology.info

Newmarket Location



Owen Sound Location



Maple Location



PATIENT NAME: _____

DATE OF BIRTH: ____/____/____ / PHONE: _____


ADDRESS: _____




HEALTH CARD #: _____ VERSION CODE _____

- CARDIOLOGISTS**
- Dr. Ali Kuny Kilany
 - Dr. S.K. Mukherjee
 - Dr. Natalie Gomperts
 - Dr. Tasnim Vira
 - Dr. Sudip Datta
 - Dr. Khaled Shelig
 - Dr. Syed Hussaini
 - Dr. Damian Redfearn
 - Dr. John Bisleri (Cardiac Surgeon St. Michael's Hospital)
 - Dr. Wael Elabbassi
 - Dr. Wael Abuzeid
- INTERNAL MEDICINE**
- Dr. Omar Rodwan
 - Dr. Abdurahman Emsalem
 - First Available

- CARDIAC TESTING**
- 2D Colour Doppler Echocardiography (ECHO) **With Contrast**
 - Treadmill Stress Echocardiogram (Stress Echo) **With Contrast**
 - Echocardiogram **With Saline "Bubble Study"** (To assess for inter-atrial shunt (PFO/ASD))
 - Graded Exercise Test (GXT)
 - Holter Monitor 48 hr. 72 hr. 7 Days 14 Days
 - Electrocardiogram (ECG)
 - Ambulatory Blood Pressure Monitor (ABPM) *Not covered by OHIP
 - CARDIOLOGY CONSULTATION**
 - URGENT (1-2 WEEKS)**
 - SEMI-URGENT (2-4 WEEKS)**
 - CONSULTATION IF ABNORMAL TEST**
- REASON FOR REFERRAL _____

- INDICATIONS**
- Palpitations/arrhythmia
 - Stroke/TIA/Cardiac Source of Embolus
 - Cardiac Murmur/Valvular heart disease
 - Shortness of breath
 - Chest pain/known coronary artery disease
 - Hypertension/Hypertensive heart disease
 - Syncope/Recurrent presyncope
 - Intermediate to high global CAD risk based on Framingham risk score
 - Significant family history of coronary artery disease
 - Assess functional capacity prior to structured exercise program (GXT)
- Other _____

- NUCLEAR CARDIOLOGY **DR. DATTA TO READ****  (Only at Maple Location)
- Myocardial Perfusion with Tc99m
 - Treadmill/Bike
 - Persantine
 - MUGA Scan with Tc99m
 - Viability Study with Thallium

- VASCULAR STUDIES (Only at Maple Location)**
- Stroke Risk Assessment** 
 - Carotid Doppler Ultrasound
 - Echocardiogram
 - Holter 72hrs Holter
 - Consultation
 - Heart failure/Edema Assessment** 
 - Chest X-ray
 - Echocardiogram
 - Peripheral Arterial Doppler
 - Peripheral Venous Doppler
 - Consultation
 - Suspected DVT/ PE assessment** 
 - Peripheral Arterial Doppler
 - Peripheral Venous Doppler
 - Echocardiogram
 - Consultation

- FOR REFERRING PHYSICIANS**
- Please send updated medications list.
 - Please send latest lab results.
 - Please send relevant cardiac testing done prior.
- FOR PATIENTS**
- Non-English speaking patients, please bring a translator to your appointment.

Send Copies To: _____

Physician's Signature: _____

Physician's Fax: _____



INSTRUCTIONS

Please arrive 10 minutes prior to your appointment time

- Bring your referral form for cardiac test.
- Bring your valid health card.
- Please provide us with 48 hours notice for cancellation.
- Please bring a complete list of your medications to all tests.

ECG

Checks electrical activity of the heart.

Duration: 5 - 10 minutes

2D Colour Doppler Echocardiography

Utilizes ultrasound waves to study the heart.

Duration: 30 - 45 minutes

Stress Exercise Test

Involves walking and/or running on the treadmill.

Please wear comfortable shoes.

Do not eat 2 hours before test.

Women: a two-piece outfit is best. No dresses.

Duration: 30 minutes

Stress Echocardiogram

Similar to exercise stress.

Addition of 2D Echo before and after exercise.

Involves walking and/or running on the treadmill.

Please wear comfortable shoes.

Women: a two-piece outfit is best. No dresses.

Duration: 1 hour

Holter Monitor

Records cardiac activity for 24, 48 or 72 hours.

No shower during the Holter is on.

Nuclear Cardiology

No caffeine for 24 hours (no coffee, tea, coke, chocolate)

No beta-blocker medication for 24 hours ex. Metoprolol, Bisoprolol, Propranolol

No erectile dysfunction medications for 72 hours

Please wear comfortable clothes and shoes suitable for exercise